



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTIONS OF EXERCISES.

284

If the angular elevation of the summits of two spires (which appear in a straight line) is α , and the angular depressions of their reflection in a lake, h feet below the point of observation, are β and γ , then the horizontal distance between the spires is

$$2h \cos^2 \alpha \sin (\beta - \gamma) \operatorname{cosec} (\beta - \alpha) \operatorname{cosec} (\gamma - \alpha).$$

[*Yale.*]

SOLUTION.

Call the point of observation P , the far summit B , the near A ; the reflection of the former C , of the latter D ; then, in PDA ,

$$\begin{aligned} PA &= PD \sin 2\beta \operatorname{cosec} (\beta - \alpha) \\ &= h \operatorname{cosec} \beta \sin 2\beta \operatorname{cosec} (\beta - \alpha) = 2h \cos \beta \operatorname{cosec} (\beta - \alpha). \end{aligned}$$

Similarly, $PB = 2h \cos \gamma \operatorname{cosec} (\gamma - \alpha)$.

$$\begin{aligned} \therefore AB &= 2h [\cos \gamma \operatorname{cosec} (\gamma - \alpha) - \cos \beta \operatorname{cosec} (\beta - \alpha)] \\ &= 2h \cos \alpha \sin (\beta - \gamma) \operatorname{cosec} (\beta - \alpha) \operatorname{cosec} (\gamma - \alpha). \end{aligned}$$

Hence the horizontal projection of AB is

$$2h \cos^2 \alpha \sin (\beta - \gamma) \operatorname{cosec} (\beta - \alpha) \operatorname{cosec} (\gamma - \alpha).$$

[*T. U. Taylor.*]

291

A regular tetrahedron and a regular octahedron are inscribed in the same sphere; compare the radii of the spheres which can be inscribed in the two solids.

[*Yale.*]

SOLUTION.

The radius of the circumscribed sphere is three-fourths the altitude of the tetrahedron and the radius of the inscribed sphere is one-fourth this altitude. Hence the radius of the sphere inscribed in the tetrahedron is $\frac{1}{3}R$, where R is the radius of the sphere. A section of the octahedron through two vertices

and the mid points of two opposite edges gives a lozenge whose diagonals are $2R$ and $R\sqrt{2}$. The perpendicular from the centre on one of these sides is the radius required and is easily found to be $\frac{1}{3}R\sqrt{3}$. The required ratio is, therefore, $1 : \sqrt{3}$. [*T. U. Taylor.*]

298

R being the radius of the circle circumscribed about a triangle, r the radius of the circle inscribed in it, and s the half sum of the sides of the triangle, the radii of the escribed circles are the roots of the equation

$$(x^2 + s^2)(x - r) = 4Rx^2.$$

[*Yale.*]

SOLUTION.

The equation may be written

$$x^3 - x^2(4R + r) + xs^2 - rs^2 = 0.$$

Call the roots r_a, r_b, r_c . It is sufficient to show that

$$\begin{aligned} r_a + r_b + r_c &= 4R + r, \\ r_ar_b + r_ar_c + r_br_c &= s^2, \\ r_ar_br_c &= rs^2, \end{aligned}$$

relations which are easily seen to be true if r_a, r_b, r_c are the radii of the escribed circles. [*T. U. Taylor.*]

EXERCISES.

308

LET points represent complex quantities in the usual way. Show that the quartic whose zeros are any four cotangential points on a fixed circular cubic, has a fixed Jacobian. [*F. Morley.*]

309

A HORIZONTAL beam of length $2a$, supported at each end, has a load in the form of an inverted parabola symmetrical with respect to the vertical line through the centre of beam. If the vertex of the parabola is b above beam, and if the load is a unit's thickness and has a heaviness unity, the deflection of the beam due to the parabolic load is $\frac{61a^4b}{360EI}$. [*T. U. Taylor.*]